

EXCLUSIVE ORAL SURGERY, LLC

'A Dental Destination Location'
Sandeep Singla DDS,MD

www.exclusiveoralsurgery.com

2055 Hamburg Turnpike
Wayne, New Jersey 07470
Tel: (973) 595-5455
Fax: (973) 595-5455



63 Valley Street
South Orange, NJ 07079
Tel:(973) 762-5773
Fax:(973) 762-5003

CONSENT FOR EXTRACTION OF TEETH AND ANESTHESIA

Patient's Name

Date

Please initial each paragraph after reading. If you have any questions, please ask your doctor BEFORE initialing.

You have the right to be informed about your diagnosis and planned surgery so that you can decide whether to have a procedure or not after knowing the risks and benefits.

Your diagnosis is: Non-Restorable tooth#

Your planned treatment is: Extraction of tooth#

Alternative treatment methods include: No Treatment/Partial Treatment/Defer Treatment.

Taking teeth out is a permanent process. Whether the procedure is easy or difficult, it is still a surgical procedure. All surgeries have some risks. They include the following and others:

1. Swelling, bruising and pain.
2. Stretching of the corners of the mouth that may lead to cracking or bruising.
3. Possible infection that might need more treatment.
4. Dry socket - jaw pain that begins a few days after surgery, that may need more care.
5. Possible damage to other teeth close to the ones being taken out, more often those with large fillings or caps.
6. Numbness, pain, or changed feelings in the teeth, gums, lip, chin and/or tongue (including possible loss of taste). This is due to the closeness of tooth roots (mainly with wisdom teeth) to the nerves which can be injured or damaged. Usually the numbness or pain goes away, but in some cases, it may need more treatment or may be permanent.

- _____ 7. Trismus – you can only open your mouth a little. This is most common after wisdom teeth are taken out. Sometimes it happens because of jaw joint (TMJ) problems already there. Damage can occur to the ligaments of the jaw joint (TMJ) from having your mouth open wide and/or for a period of time. This is more common if you already have symptoms or signs. This may need separate additional treatment.
- _____ 8. Bleeding – oozing can often happen for several hours, but a lot of bleeding is not common.
- _____ 9. Sharp ridges or bone splinters may form later at the edge of the hole where the tooth was taken out. These may need another surgery to smooth or remove.
- _____ 10. Sometimes tooth roots may be left in to avoid harming important things such as nerves or a sinus (a hollow place above your upper back teeth).
- _____ 11. The roots of the upper back teeth are often close to the sinus and sometimes a piece of root can get into the sinus. An opening may occur from the sinus into the mouth that may need more treatment.
- _____ 12. It is very rare that the jaw will break, but it is possible in cases where the teeth are buried very deep in their sockets.
- _____ 13. When donated, processed, or artificial bone substitutes are placed to preserve a socket the pieces might not join together with the natural bone and could be lost.
- _____ 14. It is understood all encounters at Exclusive Oral Surgery LLC, including my consultation/surgery/follow-up/phone calls to patients/guardians/insurance companies may be recorded for the purpose of training and/or documentation. This recording may become part of my permanent dental record or may be discarded at the sole discretion of the dental office.
- _____ 15. I have been given the opportunity to watch an informative video on extraction of wisdom teeth. This video reviews the risks/complications associated with the extraction of wisdom teeth. I fully understand these risks and agree to proceed with the proposed treatment plan of extracting the indicated teeth.

INFORMATION FOR FEMALE PATIENTS

- _____ 15. I have told my doctor that I use birth control pills. My doctor has told me that some antibiotics and other medications may reduce the preventive effect of birth control pills, and I could conceive and become pregnant. I agree to discuss with my personal doctor using other forms of birth control during my treatment, and to continue those methods until my personal doctor says that I can stop them and use only oral birth control pills.

ANESTHESIA:

LOCAL ANESTHESIA: (Novocain, Lidocaine, etc.) A shot is given to block pain in the area to be worked on.

NITROUS OXIDE WITH LOCAL ANESTHESIA: Nitrous Oxide (or Laughing Gas) helps to lessen uncomfortable sensations and offers some relaxation.

ORAL MEDICATION WITH LOCAL ANESTHESIA: A pill is taken for relaxation prior to giving local anesthesia.

INTRAVENOUS SEDATION WITH LOCAL ANESTHESIA: Makes you less aware of the procedure by making you calmer, sleepy, and less able to remember the procedure.

INTRAVENOUS GENERAL ANESTHESIA WITH LOCAL ANESTHESIA: You will be completely asleep for the procedure.

Whichever technique you choose, giving any medication involves certain risks. These include:

- ___ 1. Discomfort, swelling or bruising where the drugs are placed into a vein.
- ___ 2. Vein irritation, called phlebitis, where the drugs are placed into a vein. Sometimes this may grow to a level of discomfort or disability where it may be difficult to move your arm or hand. Sometimes medication or other treatment may be required.
- ___ 3. Nerves travel next to the blood vessels where the drugs are placed into a vein. If the needle hits a nerve or if drugs or fluid leaks out of the vessel around a nerve, you may have numbness or pain in the nerve where it runs along the arm. Usually the numbness or pain goes away, but in some cases, it may be permanent
- ___ 4. Allergic reactions (previously unknown) to any of the medications used.
- ___ 5. Nausea and vomiting, although not common, are possible unfortunate side effects. Bed rest, and sometimes medications, may be required for relief.
- ___ 6. Conscious sedation and general anesthesia are serious medical procedures and, whether given in a hospital or office, carry the risk of brain damage, stroke, heart attack or death.
- ___ 7. In situations where a breathing tube is used, you may have a sore throat, hoarseness or voice change.

YOUR OBLIGATIONS:

